

http://www.biodiversitylibrary.org/

#### Curtis's botanical magazine.

London; New York [etc.]: Academic Press [etc.] http://www.biodiversitylibrary.org/bibliography/706

#### v.56 [new ser.:v.3] (1829):

http://www.biodiversitylibrary.org/item/14216

Page(s): Title Page, Plate 2938, Text, Text

Contributed by: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 17 February 2014 5:16 PM http://www.biodiversitylibrary.org/pdf4/024554300014216

# CURTIS'S BOTANICAL MAGAZINE;

OR

## Flower Garden Displayed:

In which the most Ornamental Foreign Plants cultivated in the Open Ground, the Green-House, and the Stove, are accurately represented and coloured.

To which are added,

THEIR NAMES, CLASS, ORDER, GENERIC AND SPECIFIC CHARACTERS, ACCORDING TO THE SYSTEM OF LINNÆUS;

Their Places of Growth, Times of Flowering, and most approved Methods of Culture.

CONDUCTED

## By SAMUEL CURTIS, F. L. S.

THE DESCRIPTIONS

### By WILLIAM JACKSON HOOKER, L. L. D.

F. R. A. and L. S. and Regius Professor of Botany in the University of Glasgow.

VOL. III.

THE NEW SERIES;

Or Vol. LVI. of the whole Work.

"Soft roll your incense, Herbs, and Fruits, and Flowers, In mingled clouds, to HIM, whose sun exalts, Whose breath perfumes you, and whose pencil paints."

THOMSON.

#### LONDON:

Printed by Edward Couchman, 10, Throgmorton Street;

FOR THE PROPRIETOR, SAMUEL CURTIS,

BOTANICAL MAGAZINE WAREHOUSE, PROSPECT ROW, WALWORTH,

AND AT GLAZENWOOD, NEAR COGGESHALL, ESSEX:

Also by Sherwood, Gilbert, & Piper, 23, Paternoster Row; J. & A. Arch, Cornhill; Treuttel & Wurtz, Soho Square; Blackwood, Edinburgh; and in Holland, of Mr. Gt. Eldering, Florist, at Haarlem; And to be had of all Booksellers in Town and Country.



# CYPRIPEDIUM MACRANTHON. LARGE-FLOWERED LADY'S SLIPPER.

\*\*\*\*\*\*\*\*\*\*

Class and Order.

GYNANDRIA DIGYNIA.

(Nat. Ord.—Orchideæ.)

#### Generic Character.

Labellum ventricosum, inflatum (nunc saccatum). Columna postice terminata lobo petaloideo (stamine sterili) antheras distinguente. Petala 2 antica, sæpius connata. Br.

#### Specific Character and Synonyms.

Cypripedium \* macranthon; lobo columnæ elongato-cordato, ore labelli perianthio brevioris contracto crenulato, antheris dorso aristatis, caule folioso, foliis glabriusculis.

CYPRIPEDIUM macranthon. Swartz Gen. et Sp. Orchid. p. 103. Willd. Sp. Pl. v. 4. p. 145. Spreng. Syst. Veget. v. 3. p. 745.

Calceolus purpureus speciosus. Amman. Ruth. p. 132. n. 176. t. 21.

Calceolus y. petalis nectario æqualibus aut minoribus. Gmel. Sib. v. 1. p. 2. t. 1. y.

Descr. Stem simple, a span or more high, terete, jointed, pubescent, at the base having a sheathing scale, the root a good deal concealed by the long, sheathing bases of four or five leaves, which are ovate, attenuated at the base, wavy, striated, or almost plicated, downy at the margin and

<sup>\*</sup> From Kumpos, Venus, and modion, a slipper, Venus's Slipper, from the shape of the Labellum.

and on the nerves beneath, of a palish green colour. Peduncle terminal, enveloped by the base of a large floral leaf, glabrous. Flower solitary, large, very handsome, of an almost uniform purple colour, the two lateral or lowermost segments of the perianth, (which are united, except at the extremity, and adpressed to the underside of the labellum,) alone being greenish brown: Uppermost segment reflexed, large, broadly ovate: two inner ones broadly lanceolate, spreading, or slightly reflexed, dotted and hairy at the base within, and beautifully marked with deeper lines of purple. Labellum very large, inflated, broadly oval, striated and reticulated, the mouth contracted and crenated with a white margin: within at the base it is spotted with purple, and hairy. Column bent downward into the mouth of the labellum. Anthers large, roundish oval, deep brownishgreen, two celled, bearing on the lower part of the back a softy, fleshy spur. Abortive one, a flat, minutely glandular disk, pedunculated: from the back of which arises the elongato-cordate, petaloid lobe, of a pale reddish colour. Germen elongato-clavate, sharply angular.

This beautiful species of Cypripedium, quite new to our collections, is said by Amman to be found at Tobolsk, and by Gmelin, to be frequent in all Siberia, within the 58° of latitude, in open places, or in woods composed of scattered

Birches.

Seeds had often been sent by Dr. FISCHER of St. Petersburg to the Glasgow Botanic Garden; but we never succeeded in cultivating the plant until last year, when roots were presented to us by the same liberal Botanist. One of these, from which the present figure and description were made, blossomed under the protection of a frame in May, 1829.

C. macranthon appears to be nearly allied to the C. ventricosum, which I only know by the figure of Sweet's Brit. Fl. Garden, New Series, t. 1. But there the two innermost segments of the perianth are much narrower, and longer than the lip, the mouth of the lip is larger, and with a small cleft at the lowest extremity, and is not so regularly notched as in our plant. The whole colour too is a deeper

purple.

the later and the later to the later.

Fig. 1. Perianth, from which the Labellum (f. 2.) is removed. 3. Back view of the Column. 4. Front view of ditto. 5. Side view of ditto, slightly magnified.—Fig. 1, 2, 3, and 4, are represented of the nat. size.