

THE  
JOURNAL  
OF  
THE LINNEAN SOCIETY.

---

BOTANY.

---

---

VOL. XLI.

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.,

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1912-1913.

## ABIES SIBIRICA, Ledeb.

No. 270 a. Moist sunless hollows or shade under Pines in forests, Kushabar.

*Distribution.* The Altai Mts. and Turkestan eastwards to China, from Irkutsk province southwards to Thibet.

## LARIX SIBIRICA, Ledeb.

No. 343. Sistikem River.

No. 355. Shade of *Picea* and *Pinus* forest from 900 to 1200 m., Upper Japsa River, Sistikem. Also from the Kobdo River.

*Distribution.* The Altai Mts., Okhotsk region, and Manchuria.

## MONOCOTYLEDONES.

## ORCHIDACEÆ.

## CYTRIPEDIUM MACRANTHON, Sw.

No. 409. Natural meadows by the side of the Lower Kamsara River and under the shade of scattered Larch.

*Distribution.* S. and E. Siberia and in China.

## ORCHIS MILITARIS, Linn.

No. 368. Dry slopes of Kandart Range up to 900 m.

*Distribution.* Throughout Europe and western and southern Siberia.

## IRIDEÆ.

## IRIS ENSATA, Thunb.

No. 114. Medium dry loam in valley bottoms and shores of lakes; Achit Nor basin.

No. 178. Edges of marshes and oases in plateau desert, Kobdo Valley.

No. 475. Dry steppe association near rivers on alluvium, Dja Kul and Ulu-Kem.

*Distribution.* Dauria, Mongolia, and Afghanistan eastwards to China.

## IRIS RUTHENICA, Ait.

No. 272. Shade of *Pinus* forests, medium moist soil, Kushabar.

*Distribution.* Throughout central Asia from Transylvania to Japan.

## LILIACEÆ.

## MAIANTHEMUM BIFOLIUM, DC.

No. 317. Shade of *Pinus*, *Picea*, Birch and Poplar "Taiga," and also open glades, Anil River.

*Distribution.* Europe, N.E. Siberia, Mongolia, China, Japan, and N. America.