

## 兰科兜兰属一新种和一新变种

<sup>1</sup>陈心启 <sup>2</sup>刘仲健 <sup>2</sup>张建勇

<sup>1</sup>(中国科学院植物研究所系统与进化植物学开放研究实验室 北京 100093)

<sup>2</sup>(深圳市梧桐山苗圃总场 深圳 518114)

### A new species and a new variety of *Paphiopedilum* (Orchidaceae)

<sup>1</sup>CHEN Sing-Chi <sup>2</sup>LIU Zhong-Jian <sup>2</sup>ZHANG Jian-Yong

<sup>1</sup>(*Laboratory of Systematic and Evolutionary Botany, Institute of Botany, the Chinese Academy of Sciences, Beijing 100093*)

<sup>2</sup>(*Shenzhen City Wutongshan Nurseries, Shenzhen 518114*)

**Key words** *Paphiopedilum*; *P. tridentatum*; *P. armeniacum* var. *undulatum*; New species; New variety; Vietnam; China

**关键词** 兜兰属; 齿瓣兜兰; 浅黄兜兰(变种); 新种; 新变种; 越南; 中国

#### 齿瓣兜兰 新种 图 1

***Paphiopedilum tridentatum* S. C. Chen et Z. J. Liu, sp. nov.** Fig. 1

Species nova *Paphiopedilo appletoniano* (Gower) Rolfe similis, a quo bene differt sepalo intermedio subplano ad marginem basis non recurvato, petalis apice tridentatis, staminodio purpureo-nigro praeter marginem et centrum viridulum apice mucronato et sine sinus.

Rhizoma breve. Folia 4~5, disticha, anguste elliptico-oblonga, 10~16 cm longa, 1.8~3.5 cm lata, supra pallido-viridi- et atroviridi-tessellata, infra pallide viridia et ad basin plus minusve purpurea, carinata, apice breviter acuminata, basi in petiolum 1~2 cm longum conduplicatum contracta. Scapus suberectus, ca. 45 cm longus, purpureo-chocolatinus, dense albo-pubescent, uniflorus; bractea late ovata, ca. 1.4 cm longa, 1.2 cm lata, plus minusve conduplicata, viridula, ad basin purpureo-chocolatina, extus albo-pubescent, apice leviter tridentata; pedicellus cum ovario ca. 6 cm longus, purpureo-chocolatinus, dense albo-pubescent; flos ca. 10 cm in diametro; sepalum intermedium late ovatum, subplanum, ca. 3.2 cm longum, 2.5 cm latum, viride, in dimidio inferiore longitudinaliter purpureo-nervosum, prope basin transverse chocolatino-fasciarium, apice subacutum, dorso albo-puberulum, minute ciliatum; synsepalam ovatum, concavum, ca. 2.6 cm longum, 1.6 cm latum, viride, obscure longitudinaliter purpureo-nervosum, dorso albo-puberulum, minute ciliatum; patala spathulata, ca. 5 cm longa, 2.1 cm lata, in dimidio superiore purpurella, infra medium viridi-purpurella et nigro-punctata, apice subrotundata et tridentata, basi ventrali villosa, minute ciliata; labellum galeiformi-saccatum, ca. 4.6 cm longum, viridi-purpurellum, sacco obovato-oblongo ca. 2.4 cm longo 1.8 cm lato extus glabro ad basin interiores villosa, ungue ca. 2.2 cm longo 1.2 cm lato, lobis lateralibus incurvis verruculatis; staminodium fere transverse elliptico-oblongum, ad basin leviter angustatum, ca. 8 mm longum, 10 mm latum, apice mucronatum, basi auriculatum, purpureo-nigrum praeter marginem et centrum viridulum.

**Vietnam** (越南). without precise locality (无详细地点), cult. plant, 2001-03-28, Z. J. Liu 20141 (holotype, here designated, in Herbarium, Shenzhen City Wutongshan Nurseries).




图1 齿瓣兜兰 1. 带花植株; 2. 中萼片, 正面观; 3. 合萼片, 正面观; 4. 脣瓣, 侧面观; 5. 蕊柱, 侧面观; 6. 退化雄蕊, 正面观。

Fig. 1 *Paphiopedilum tridentatum* S. C. Chen et Z. J. Liu 1. Flowering plant; 2. Dorsal sepal, front view; 3. Synsepal, front view; 4. Lip, side view; 5. Column, side view; 6. Staminode, front view.




图2 浅黄兜兰(变种) 1. 带花植株; 2. 中萼片, 正面观; 3. 合萼片, 正面观; 4. 花瓣, 正面观; 5. 蕊柱, 侧面观; 6. 退化雄蕊, 正面观; 7. 子房, 横切面。

Fig. 2 *Paphiopedilum armeniacum* S. C. Chen et F. Y. Liu var. *undulatum* Z. J. Liu et J. Y. Zhang 1. Flowering plant; 2. Dorsal sepal, front view; 3. Synsepal, front view; 4. Petal, front view; 5. Column, side view; 6. Staminode, front view.; 7. Ovary cross section.

根状茎短。叶 4~5 枚,二列,狭椭圆状矩圆形,长 10~26 cm,宽 1.8~3.5 cm,上面具有深浅绿色相间的网格斑,背面浅绿色并在基部多少具紫晕,有龙骨状突起,先端短渐尖,基部收狭成柄;柄长 1~2 cm,对褶。花葶近直立,长约 45 cm,紫栗色,密被白色短柔毛,具 1 花;花苞片宽卵形,长约 1.4 cm,宽 1.2 cm,多少对褶,绿色,基部紫栗色,背面被白色短柔毛,先端稍具 3 小齿;花梗和子房长约 6 cm,紫栗色,密被白色短柔毛;花直径约 10 cm;中萼片宽卵形,近平坦,长约 3.2 cm,宽约 2.5 cm,绿色,下半部具浅紫色纵脉,近基部具一条横的栗色带,先端近急尖,背面有白色细柔毛,边缘具细缘毛;合萼片卵形,凹陷,长约 2.6 cm,宽 1.6 cm,绿色,有模糊的淡紫色纵脉,背面被白色细柔毛,边缘具细缘毛;花瓣匙形,长约 5 cm,宽 2.1 cm,淡紫色,中部以下为带绿的淡紫色并有黑色斑点,先端近浑圆并有 3 小齿,内面基部有长柔毛,边缘具细缘毛;唇瓣盔状囊形,长约 4.6 cm,带绿的淡紫色;囊倒卵状矩圆形,长约 2.4 cm,宽 1.8 cm,外面无毛,内面基部有长柔毛;柄长约 2.2 cm,宽 1.2 cm,内弯的侧裂片上有小疣状突起;退化雄蕊近横椭圆状矩圆形,基部稍狭,长约 8 mm,宽 10 mm,先端具短尖头,基部有耳,紫黑色,但边缘与中央淡绿色。

此新种与卷萼兜兰 *Paphiopedilum appletonianum* 有亲缘关系,但本新种中萼片近平坦,基部两侧边缘不反卷;花瓣先端具 3 小齿;退化雄蕊除边缘与中央外均紫黑色,先端具短尖头,不具弯缺,易于分辨。

### 浅黄兜兰 新变种 图 2

***Paphiopedilum armeniacum* S. C. Chen et F. Y. Liu var. *undulatum* Z. J. Liu et J. Y. Zhang, var. nov. Fig. 2**

A typo differt floribus plerumque duobus minoribus 4.5~5 cm in diametro flavidis, petalis 1.4~1.7 cm latis ad maginem valde undulatis.

**China. Western Yunnan** (云南西部); without precise locality(无详细地点), cult. plant, 2001-03-17. Z. J. Liu 21317 (holotype, here designated, in Herbarium, Shenzhen City Wutongshan Nurseries).

新变种区别于原变种在于:花通常 2 朵,较小,直径 4.5~5 cm,淡黄色;花瓣宽 1.3~1.7 cm,边缘明显波状。

In the present paper two new taxa, *Paphiopedilum tridentatum* S. C. Chen et Z. J. Liu and *Paphiopedilum armeniacum* S. C. Chen et F. Y. Liu var. *undulatum* Z. J. Liu et J. Y. Zhang, are described and diagnosed respectively. *Paph. tridentatum* is a Vietnamese species, very similar to *Paph. appletonianum* (Gower) Rolfe at first glance, but their floral structure is quite different from each other. The new species has a nearly flat dorsal sepal, tridentate-tipped petals and an almost purple-black staminode with mucronate apex, while *Paph. appletonianum* possesses a dorsal sepal strongly recurved along its basal margins, edentate petals and a greenish staminode with a clear sinus at its apex. *Paph. armeniacum* var. *undulatum* is said to be collected from western Yunnan. It differs from *Paph. armeniacum* var. *armeniacum* by having usually two yellowish flowers 4.5~5 cm across, and strongly undulate-margined petals 1.4~1.7 cm wide. Thw two new taxa are both cultivated plants, which were transplanted from a nursery in southwestern Yunnan.