

http://www.biodiversitylibrary.org/

Curtis's botanical magazine.

London; New York [etc.]: Academic Press [etc.] http://www.biodiversitylibrary.org/bibliography/706

v.47 (1819-1820): http://www.biodiversitylibrary.org/item/14331

Page(s): Title Page, Plate 2129, Text, Text

Contributed by: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 17 February 2014 5:15 PM http://www.biodiversitylibrary.org/pdf4/024554200014331

CURTIS'S Botanical Magazine;

OR,

FLOWER-GARDEN DISPLAYED:

IN WHICH

The most Ornamental Foreign Plants, cultivated in the Open Ground, the Green-House, and the Stove, are accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according to the celebrated Linnæus; their Places of Growth, and Times of Flowering;

Together with the most approved Methods of CULTURE.

AWORK

Intended for the Use of such Ladies, Gentlemen, and Gardeners, as wish to become scientifically acquainted with the Plants they cultivate.

By JOHN SIMS, M.D.

FELLOW OF THE ROYAL AND LINNEAN SOCIETIES.

VOL XLVII.

Being the Fifth of the New Series.

The Flowers, which grace their native beds,
Awhile put forth their blushing heads,
But, e'er the close of parting day,
They wither, shrink, and die away:
But these, which mimic skill hath made,
Nor scorched by suns, nor killed by shade,
Shall blush with less inconstant hue,
Which ART at pleasure can renew.

Lloyd.

London;

Printed by W. & S. Couchman, Throgmorton-Street.

Published by Sherwood, Neely, & Jones, 20, Paternoster-Row;

And Sold by the principal Booksellers in Great-Britain and Ireland.

M DCCCXX.



CYPRIPEDIUM VENUSTUM. COMELY LADY'S SLIPPER.

becoming quite upright, and the lateral ones spreading hori-

Class and Order.

GYNANDRIA DIGYNIA:

Generic Character.

plant, is a mere notice by the little in a letter to

Labellum (nectarium) ventricosum, inflatum (nunc saccatum). Columna postice terminata lobo petaloideo (stamine sterili) antheras distinguente. Petala 2 antica sæpius connata. Brown in Hort. Kew.

Specific Character and Synonym.

Cypripedium venustum; acaule, foliis lanceolatis equitantibus maculatis, petalis lateralibus ciliatis, nectario venoso ore inflexo.

Cypripedium venustum. Wallich.

DESCR. Leaves ovate-lanceolate, carinate, spreading distichwise, their bases overlapping one another alternately; spotted with large green spots on the upper surface, and small purple dots on the under, perennial. Scape but little longer than the leaves, rounded. Spathe ovate, concave, keeled; keel and margins blackish. Germen twice as long as the spathe, obtusely 3-angled. Corolla at first nodding: upper petal oval, concave, white, striped with green: lower petal smaller, otherwise similar, composed of two united, which is proved (as observed by Brown) by the absence of the midrib. The two lateral petals longer than the others, strap-shaped, ciliated with dark purple hairs, externally of the same colour with the others; internally marked with black round spots, and towards the tip tinged with purple. Slipper (labellum or nectary) inflated, oval, beautifully veined with green and tinged with purple, the opening bent inward at the sides, leaving the corners erect. Column short, dark green, terminated with a twolobed lid. Style incurved, short; stigma large, orbicular. Filaments Filaments lateral, sulcate; anthers at the time of examination

much wasted: pollen glutinous.

The Germen, which was much incurved at the time the drawing was taken, afterwards became straighter, and the flower in consequence ceased to droop; the upper petal becoming quite upright, and the lateral ones spreading hori-

zontally.

Native of the East Indies. Communicated by Messrs. Whitley, Brames, and Milne, in November; who received it from the botanic garden at Calcutta. It was brought here by Captain Craigie. The only account we can find of this plant, is a mere notice by Dr. Wallich in a letter to Dr. Francis Hamilton, published in the first volume of the Edinburgh Philosophical Journal, of a third new species of Cypripedium from Nepal, to which, we understand from Mr. Brown, the Doctor has given the name, which we have adopted.

And the contract of the first the contract.

Organications ventustum recuie, folis danceolatis equitan.

oreliniexo.

tibus maculatis, petalis lateralibus dilatis, nectario venose

elgue, leite bes erelges sympoth to eloce as a diw

wise; their bases orgalispoing one another alternated.; mosted

the main particle of the first and acceptable fair and a political than the first and the first and

leages, rounded. Spring wrate, concave, Lealed; host and mars

A PRINCIPAL CONTRACTOR DELL'ARTER DE L'ARTER LES BERRES DE L'ARTER DE L'ARTER

Sangled wells at the bodding a sent bound of the contract of t

cave, white, stained with execut; locality lights, out to aite

devised as the property of the search of the

by Brown) by the absence of the midden. The two two taterof.

petats longer than the others strangard, ciliated with dark

purple bairs, externally of the same colour with the others;

internally marked with black round spots, and towards the

up tinged with purple. Stipper (inheflum or mentary) inficied;

oval, beautifully veliced with green and tinged with purple,

the opening bent inggard at the sides, lenging the corners

erect, Cobsess short, dark green, terminated with a two-

lobed lid. 'Style incurred, short; sligma large, orbicular.